BILLING AND CODING GUIDE



PATIENT COVERAGE

- 1) Complete a **TROGARZO[®] Enrollment Form** and FAX the patient's enrollment package to THERA patient support[™] at 1-855-836-3069.
- 2) A Patient Care Coordinator will then assess and advise on the patient's private or government insurance coverage, including AIDS Drug Assistance Programs (ADAPs). The Patient Care Coordinator will also assist in applying any eligible co-pay assistance.
- 3) TROGARZO[®] may then be ordered and given at your office.

HOW TO ORDER

TROGARZO® Authorized Distributor: CuraScript SD Acute: 800.211.1455

TELEPHONE	1-877-599-7748 (Mon-Fri 8:30AM-7PM ET)	
FAX	1-800-862-6208	
WEBSITE	www.curascriptsd.com	

For information on payment, shipping or return policies, please contact Curascript SD directly.



Questions?

Contact us at 1-833-23-THERA (1-833-238-4372), Mon-Fri 8:30AM-8PM ET

See back page for full Important Safety Information.

Note: Individual payer organizations should be contacted for coverage and reimbursement policies and processes, including prior authorization, if necessary.

PRODUCT INFORMATION

TROGARZO® is available in a carton containing two single-dose vials.



Each vial contains 200 mg ibalizumab-uiyk

TROGARZO® DOSE	VIALS REQUIRED	CARTONS REQUIRED
Loading Dose (2,000 mg ibalizumab-uiyk)	10	5
Maintenance Doses (800 mg ibalizumab-uiyk)	4	2

NDC	DESCRIPTION	
62064-122-02	Pack of 2 vials, each containing 200 mg of ibalizumab-uiyk for intravenous use	
ICD-10 CODE	DESCRIPTION	
B20	Human immunodeficiency virus (HIV) disease	
Z16.33	Resistance to antiviral drug(s)	
CPT CODE	DESCRIPTION	
96365	IV infusion for therapy, prophylaxis, or diagnosis; initial, up to 1 hour	
96374	IV push, single or initial substance/drug lasting 15 minutes or less	

HCPCS CODE	DESCRIPTION
J1746	Injection, ibalizumab-uiyk, 10 mg

Please check with payor to verify coding or special billing requirements. Correct coding is the responsibility of the provider submitting a claim for the item or service.

See back page for full Important Safety Information.

SAMPLE CMS-1500 CLAIM FORM

PICA		PICA 🗌
1. MEDICARE MEDICAID TRICARE CHAMP		(For Program in Item 1)
(Medicare#) (Medicaid#) (ID#/DoD#) (Member 2. PATIENT'S NAME (Last Name, First Name, Middle Initial)	(<i>ID#</i>) (<i>ID#</i>) (<i>ID#</i>) PATIENT'S BIRTH DATE SEX 4. INSURED'S NAME (Last Na MM + DD + YY SEX	me, First Name, Middle Initial)
5. PATIENT'S ADDRESS (No., Street)	PATIENT RELATIONSHIP TO INSURED 7. INSURED'S ADDRESS (No Self Spouse Child Other	, Street)
CITY STATE	RESERVED FOR NUCC USE CITY	STATE
ZIP CODE TELEPHONE (Include Area Code)	ZIP CODE	TELEPHONE (Include Area Code)
OTHER INSURED'S NAME (Last Name, First Name, Middle Initial)	, IS PATIENT'S CONDITION RELATED TO: 11. INSURED'S POLICY GRO	
. OTHER INSURED'S POLICY OR GROUP NUMBER	EMPLOYMENT? (Current or Previous) a. INSURED'S DATE OF BIRT MM DD Y YES NO I	H SEX
RESERVED FOR NUCC USE	AUTO ACCIDENT? PLACE (State) b. OTHER CLAIM ID (Designa	
RESERVED FOR NUCC USE	OTHER ACCIDENT?	DR PROGRAM NAME
	YES NO	
I. INSURANCE PLAN NAME OR PROGRAM NAME	bd. CLAIM CODES (Designated by NUCC)	TH BENEFIT PLAN? If yes, complete items 9, 9a, and 9d.
READ BACK OF FORM BEFORE COMPLETIN 2. PATIENT'S OR AUTHORIZED PERSON'S SIGNATURE I authorize th	ase of any medical or other information necessary payment of medical benefit	ZED PERSON'S SIGNATURE I authorize s to the undersigned physician or supplier fo
to process this claim. I also request payment of government benefits eithe below.	vyself or to the party who accepts assignment services described below.	
SIGNED	ATESIGNED	
MM DD YY QUAL	MM DD YY MM DD FROM	TO WORK IN CURRENT OCCUPATION YY MM DD YY TO D
7. NAME OF REFERRING PROVIDER OR OTHER SOURCE	I8. HOSPITALIZATION DATE: MM DD FROM I	S RELATED TO CURRENT SERVICES YY MM DD YY TO I
9. A 21 LAIM INFORMATION (Designated by NUCC)	20. OUTSIDE LAB?	\$ CHARGES
1. DIAG. OR NATURE OF ILLNESS OR INJURY Relate A-L to se	YES NO Ine below (24E) ICD Ind. 22. RESUBMISSION CODE	ORIGINAL REF. NO.
A B20 B. Z16.33 c.	D 23, PRIOR AUTHORIZATION	
F G	H. L 20. Thior comonization	NOMBER
AA DATE(S) OF SERVICE B. C. C rom To PLACE OF DD YY MM DD YY SERVICE EMG	ES, SERVICES, OR SUPPLIES usual Circumstances) MODIFIER MODIFIER MODIFIER S CHARGES 24	G H. I. J. SDT ID. RENDERING Han QUAL. PROVIDER ID. #
62064-122-02, TROGARZO INJ		
j j j J17	80	NPI
963	1	NPI
oi 963	1	NPI
		NPI
		NPI
		NPI
25. FEDERAL TAX I.D. NUMBER SSN EIN 26. PATIENT'S	OUNT NO. 27. ACCEPT ASSIGNMENT? 28. TOTAL CHARGE 2000 (Corgov. claims, see back) Yes NO \$	29. AMOUNT PAID 30. Rsvd for NUCC

21 Inter app

Enter applicable ICD-10 Diagnosis Codes:

• B20

• Z16.33



Enter product information (NDC) with qualifier:

• 62064-122-02



Enter Drug Code (HCPCS) and Procedure Code (CPT):

- J1746
- 96365 or 96374

24G

For TROGARZO[®], enter billing units according to the correct dose:

- For a **Loading Dose** (2,000 mg ibalizumab-uiyk): Enter 200
- For **Maintenance Doses** (800 mg ibalizumab-uiyk): Enter 80

Sample provided for reference only. Theratechnologies Inc. makes no guarantee of coverage by payors. Please check with payor to verify coding or special billing requirements. Correct coding is the responsibility of the provider submitting a claim for the item or service.

IMPORTANT SAFETY INFORMATION

Indication

TROGARZO[®] (ibalizumab-uiyk), in combination with other antiretroviral(s), is indicated for the treatment of human immunodeficiency virus type 1 (HIV-1) infection in heavily treatment-experienced adults with multidrug resistant HIV-1 infection failing their current antiretroviral regimen.

Contraindications

TROGARZO[®] is contraindicated in patients with a prior hypersensitivity reaction to TROGARZO[®] or any components of the product.

Warnings and Precautions

Hypersensitivity Including Infusion-Related and Anaphylactic Reactions

Hypersensitivity reactions including infusion-related reactions and anaphylactic reactions have been
reported following infusion of TROGARZO[®] during post-approval use. Symptoms may include dyspnea,
angioedema, wheezing, chest pain, chest tightness, cough, hot flush, nausea, and vomiting. If signs and
symptoms of an anaphylactic or other clinically significant hypersensitivity reaction occur, immediately
discontinue administration of TROGARZO[®] and initiate appropriate treatment. The use of TROGARZO[®] is
contraindicated in patients with known hypersensitivity with TROGARZO[®].

Immune Reconstitution Inflammatory Syndrome

• Immune Reconstitution Inflammatory Syndrome (IRIS) has been reported in one patient treated with TROGARZO[®] in combination with other antiretrovirals. During the initial phase of combination antiretroviral therapies, patients whose immune systems respond may develop an inflammatory response to indolent or residual opportunistic infections, which may necessitate further evaluation and treatment.

Embryo-Fetal Toxicity

 Based on animal data, TROGARZO[®] may cause reversible immunosuppression (CD4+ T cell and B cell lymphocytopenia) in infants born to mothers exposed to TROGARZO[®] during pregnancy. Immune phenotyping of the peripheral blood and expert consultation are recommended to provide guidance regarding monitoring and management of exposed infants based on the degree of immunosuppression observed. The safety of administering live or live-attenuated vaccines in exposed infants is unknown.

Adverse Reactions

The most common adverse reactions (all Grades) seen in clinical trial experience, reported in at least 5% of subjects receiving TROGARZO® were diarrhea (8%), dizziness (8%), nausea (5%) and rash (5%). Most (90%) of the adverse reactions reported were mild or moderate in severity. Two subjects experienced severe adverse reactions: one subject had a severe rash and one subject developed IRIS manifested as an exacerbation of progressive multifocal leukoencephalopathy.

Use in Specific Populations

- **Pregnancy:** No adequate human data are available to establish whether or not TROGARZO[®] poses a risk to pregnancy outcomes. Monoclonal antibodies, such as ibalizumab-uiyk, are transported across the placenta as pregnancy progresses; therefore, ibalizumab-uiyk has the potential to be transmitted from the mother to the developing fetus.
- Lactation: No data are available regarding the presence of TROGARZO[®] in human milk, the effects on the breastfed child, or the effects on milk production. Because of the potential for HIV-1 transmission, instruct mothers not to breastfeed if they are receiving TROGARZO[®].

Please see enclosed full Prescribing Information for TROGARZO®.

To report suspected adverse reactions, contact THERA patient support® (1-833-238-4372) or the FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.



