

MAKE UNDETECTABLE POSSIBLE

Take Action

with TROGARZO®
(ibalizumab-uiyk)



**A CONVENIENT
30- SECOND IV PUSH
ADMINISTRATION OPTION†**

† After the 30-minute IV infusion loading dose, all subsequent doses can be given in 30 seconds every 2 weeks with the IV push

You can make undetectable possible and may lower your pill count. Ask your healthcare provider about TROGARZO® (ibalizumab-uiyk).

TROGARZO® (ibalizumab-uiyk) is a prescription medicine that is used with other antiretroviral medicines to treat Human Immunodeficiency Virus-1 (HIV-1) infection in adults who:

- have received anti-HIV-1 regimens in the past, and
- have HIV-1 virus that is resistant to many antiretroviral medicines, and
- who are failing their current antiretroviral therapy

IMPORTANT SAFETY INFORMATION

Do not receive TROGARZO® if you have had an allergic reaction to TROGARZO® or any of the ingredients in TROGARZO®.

Please see Important Safety Information throughout and the full Prescribing Information for TROGARZO® online at www.trogarzo.com.




Trogarzo®

(ibalizumab-uiyk)
Injection

200 mg/1.33 mL (150 mg/mL)

WHAT IS TROGARZO®?

TROGARZO® is a fully active medicine that's used in combination with other antiretroviral medicines (ARV) to treat HIV-1 infection in adults who:



UNDETECTABLE:* A POSSIBILITY AGAIN

If you have been on different HIV-1 treatments in the past and your current ARV regimen is failing, you deserve to know about all of your treatment options, including TROGARZO® - an HIV-1 treatment specifically designed for people like you.

Together, you and your healthcare provider can decide if TROGARZO® should be part of your treatment regimen to help keep your viral load in check.

* Undetectable viral load is defined as fewer than 50 copies of HIV RNA per mL of blood.



**ASK YOUR HEALTHCARE PROVIDER
IF TROGARZO® IS RIGHT FOR YOU**

IMPORTANT SAFETY INFORMATION

TROGARZO® can cause serious side effects, including:

- **Allergic reactions.** TROGARZO® can cause allergic reactions, including serious reactions, during and after infusion. Tell your healthcare provider or nurse, or get medical help right away if you get any of the following symptoms of an allergic reaction: trouble breathing, swelling in your throat, wheezing, chest pain, chest tightness, cough, hot flush, nausea or vomiting.
- **Changes in your immune system (Immune Reconstitution Inflammatory Syndrome)** can happen when you start taking HIV-1 medicines. Your immune system might get stronger and begin to fight infections that have been hidden in your body for a long time. Tell your healthcare provider right away if you start having new symptoms after receiving TROGARZO®.

Please see Important Safety Information throughout and the enclosed full Prescribing Information for TROGARZO®. The TROGARZO® Prescribing Information is also available online at www.trogarzo.com.

UNDETECTABLE MAY BE WITHIN REACH.

CAN TROGARZO® REDUCE MY VIRAL LOAD?

Did you know that if your virus becomes resistant to multiple HIV-1 medications, treatment guidelines suggest considering using a drug that works differently than others you may have tried?

TROGARZO® works differently. TROGARZO® binds to the T-cell's surface to block HIV-1 from entering. This protects your immune cells from the virus.

THIS CAN REDUCE YOUR VIRAL LOAD.

In a clinical study, TROGARZO® was proven to have a significant and long-lasting impact on the amount of HIV-1 in the body, reducing the viral load.

83%

OF PATIENTS EXPERIENCED A **DECREASE IN THEIR VIRAL LOAD** OF AT LEAST 66% ONE WEEK AFTER RECEIVING THEIR LOADING DOSE OF TROGARZO®.

Studies have also shown that **50% of heavily treatment-experienced patients are able to achieve and maintain viral suppression (<200 copies/mL) through 25 weeks of treatment with TROGARZO®.**

30 SECONDS EVERY 2 WEEKS VIA IV PUSH

Did you know that there is another option to taking daily pills?

TROGARZO® is not a daily pill. Once you've received your initial loading dose by intravenous (IV) infusion, all subsequent doses can be given **in 30 seconds** every 2 weeks via IV push.



Given intravenously **every 2 weeks**



Appointments arranged by your personal **Patient Care Coordinator**



Given by a **healthcare provider** at home, an infusion center or healthcare provider's office

WHAT ARE THE BENEFITS OF A 2-WEEK TOUCHPOINT?

The TROGARZO® administration schedule means that throughout your TROGARZO® journey you get care and support. You can take this opportunity to ask questions about your health or treatment.

IMPORTANT SAFETY INFORMATION

Before you receive TROGARZO® (ibalizumab-uiyk), tell your healthcare provider about all of your medical conditions, including if you are:

- **Pregnant or plan to become pregnant.** It is not known if TROGARZO® may harm your unborn baby. Tell your healthcare provider if you become pregnant during treatment with TROGARZO®.

Also tell your healthcare provider about all the medicines you take, including all prescription and over-the-counter medicines, vitamins, and herbal supplements.

Please see Important Safety Information throughout and the enclosed full Prescribing Information for TROGARZO®. The TROGARZO® Prescribing Information is also available online at www.trogarzo.com.

93% OF TROGARZO® PATIENTS SURVEYED RECOMMEND TROGARZO® TO OTHER APPROPRIATE PATIENTS.

87%

REPORTED THAT HAVING A **2-WEEK TOUCHPOINT** WITH AN HCP WAS A POSITIVE EXPERIENCE.

80%

REPORTED THAT HAVING A **2-WEEK TOUCHPOINT** WITH AN HCP HELPED THEM ADHERE TO THEIR OTHER HIV MEDICATIONS.


[†] This single-blinded study required patients (n=30) to complete a brief online survey regarding patient experience with TROGARZO®. Respondents had been taking TROGARZO® an average of 8.5 months and average duration of treatment ranged from 1 to 18 months.* The survey included 20 questions of various types (i.e., Likert scale, dichotomous, multiple choice, open-ended), which were analyzed in aggregate and by infusion setting (i.e., HCP clinic, infusion center, or at home). All surveys were completed from December 2020 - March 2021 (prior to the availability of the IV push administration method), and all participants were compensated for their time.

[‡] Applies to 15/30 respondents. Information for the remaining 15 respondents was not available.

TROGARZO[®], A POWERFUL HIV-1 TREATMENT

Given as an initial loading dose as an infusion, TROGARZO[®] can then **be given once every 2 weeks via IV push**. TROGARZO[®] was the first available long-acting antiretroviral against HIV-1 – a breakthrough in a class of its own.

How does the dosing of TROGARZO[®] make it different from pills?



If the level of medication in your body falls into the “red zone”, it means that the level may not be high enough to suppress your viral load and your virus can become resistant to your current medication.

TROGARZO[®] is long-acting, which means that it maintains a sufficient level of medication in the body over its dosing period of 14 days

A PROVEN TOLERABILITY & SAFETY PROFILE



TROGARZO® is generally well tolerated.

The most common side effects of TROGARZO® were diarrhea, dizziness, nausea, and rash. Most adverse reactions reported were mild to moderate in severity.



Did you know that some HIV-1 medications can cause interactions with other drugs which can reduce their efficacy or increase side effects?

TROGARZO® does not interact with other drugs, so you can take it with other medications you may be taking. Be sure to discuss other medications you may be taking with your healthcare provider before taking TROGARZO®.

IMPORTANT SAFETY INFORMATION

Before you receive TROGARZO® (ibalizumab-uiyk), tell your healthcare provider about all of your medical conditions, including if you are:

- **Breastfeeding or plan to breastfeed.** You should not breastfeed if you have HIV-1 because of the risk of passing HIV-1 to your baby. Do not breastfeed if you are receiving TROGARZO® as it is not known if TROGARZO® passes into breast milk. Talk with your healthcare provider about the best way to feed your baby during treatment with TROGARZO®.

The most common side effects of TROGARZO® include diarrhea, dizziness, nausea, and rash. These are not all the possible side effects of TROGARZO®.

Do not receive TROGARZO® if you have had an allergic reaction to TROGARZO® or any of the ingredients in TROGARZO®.

Please see the back cover for complete safety details.

WHERE CAN I HAVE TROGARZO[®] ADMINISTERED?



**INFUSION
CENTER**



YOUR HOME



**YOUR HEALTHCARE
PROVIDER'S
OFFICE**

Depending on your insurance coverage, your Patient Care Coordinator will arrange for your treatment to be given at a location of your choosing - at an infusion center, in your own home, or at your healthcare provider's office. If you decide over time that another location would be more convenient for you, simply let them know.

You can also reach them by phone to ask any questions you may have about your treatment.

WHAT TO EXPECT DURING TROGARZO® ADMINISTRATION

Before

- Ask your Patient Care Coordinator if you should bring any documents (like insurance information) to your appointment.


During

- Someone will monitor you during treatment.
- Take the opportunity to discuss HIV-related topics with the care provider.

After

- You'll also be monitored afterwards for a short period of time to ensure that your body tolerates your treatment well.

Please see the back cover for complete safety details.



“Know your options and get all the information. Do you want to continue the course you’re on or try something new? I’ve seen my viral load and CD4 improve, as well as my overall health.”

-Jason, a real TROGARZO® patient



SCAN HERE TO HEAR ABOUT THE EXPERIENCE OF JASON, A REAL TROGARZO® PATIENT

This is the experience of one patient and may not be representative of all TROGARZO® patients.

MEET THERA PATIENT SUPPORT®

We're here to help make your journey as smooth as possible and answer any questions you may have.

Once TROGARZO® becomes a part of your regimen, you'll receive a welcome call from your dedicated Patient Care Coordinator.

Thinking about how to afford your treatment?

Don't worry, TROGARZO® is widely covered by all commercial and government insurance plans, including ADAP. Commercially insured patients may be eligible for co-pay assistance and may pay as little as \$0.



FROM APPOINTMENT REMINDERS TO INSURANCE COVERAGE ASSISTANCE, AND ANYTHING IN BETWEEN, **THERA PATIENT SUPPORT® IS HERE TO HELP.**

LEARN MORE AT [TROGARZO.COM](https://trogarzo.com)

 **THERA** patient support®

IS TROGARZO® RIGHT FOR YOU?

START THE CONVERSATION

Take this quiz to determine if TROGARZO® might be right for you:

1. Are you having any challenges taking your current HIV medications?	<input type="checkbox"/> YES <input type="checkbox"/> NO
2. Do you skip or miss your HIV medication doses sometimes?	<input type="checkbox"/> YES <input type="checkbox"/> NO
3. Have you changed your HIV medications in the past?	<input type="checkbox"/> YES <input type="checkbox"/> NO
4. Are you taking your HIV medication and your viral load is still detectable?	<input type="checkbox"/> YES <input type="checkbox"/> NO
5. Would you be open to including something other than a daily pill in your treatment regimen if given the choice?	<input type="checkbox"/> YES <input type="checkbox"/> NO
6. Do you consider safety an important part of your next HIV medication choice?	<input type="checkbox"/> YES <input type="checkbox"/> NO

If you answered “YES” to 4 out of 6 questions, you might be a potential TROGARZO® candidate.

**At your next visit with your healthcare provider,
ask to learn more about TROGARZO®!**

If your current HIV-1 treatment is failing, it is important to know all of your treatment options. Ask your healthcare provider if TROGARZO® should be part of your regimen to keep your viral load in check.

Because you deserve to know.

FOR MORE INFORMATION, VISIT TROGARZO.COM

You can view and download the discussion guide from the website! Remember to bring a copy with you to your next doctor’s appointment.

Please see the next page for complete safety details.

SO, IS TROGARZO® RIGHT FOR YOU?



**A CONVENIENT
30- SECOND IV PUSH
ADMINISTRATION OPTION†**

† After the 30-minute IV infusion loading dose, all subsequent doses can be given in 30 seconds every 2 weeks with the IV push

Have you changed ARV regimens multiple times in the past? Are you looking for a different option because your current regimen is no longer working?

LEARN MORE AT TROGARZO.COM

 **THERA patient support®**

If you have any questions, call the THERA patient support® program at **1-833-23 THERA (1-833-238-4372)**,
Monday-Friday, 8:30AM-8PM ET.

IMPORTANT SAFETY INFORMATION

TROGARZO® (ibalizumab-uiyk) is a prescription medicine that is used with other antiretroviral medicines to treat Human Immunodeficiency Virus-1 (HIV-1) infection in adults who:

- have received anti-HIV-1 regimens in the past, and
- have HIV-1 virus that is resistant to many antiretroviral medicines, and
- who are failing their current antiretroviral therapy

It is not known if TROGARZO® is safe and effective in children.

TROGARZO® can cause serious side effects, including:

- Allergic reactions. TROGARZO® can cause allergic reactions, including serious reactions, during and after infusion. Tell your healthcare provider or nurse, or get medical help right away if you get any of the following symptoms of an allergic reaction: trouble breathing, swelling in your throat, wheezing, chest pain, chest tightness, cough, hot flush, nausea or vomiting.
- Changes in your immune system (Immune Reconstitution Inflammatory Syndrome) can happen when you start taking HIV-1 medicines. Your immune system might get stronger and begin to fight infections that have been hidden in your body for a long time. Tell your healthcare provider right away if you start having new symptoms after receiving TROGARZO®.

For more information or medical advice about side effects, ask your healthcare provider. You may report side effects to the FDA at 1-800-FDA-1088 or the THERA patient support® program at 1-833-238-4372.

Please see Important Safety Information throughout and the enclosed full Prescribing Information for TROGARZO®. The TROGARZO® Prescribing Information is also available online at www.trogarzo.com.



TROGARZO® is a registered trademark of TaiMed Biologics Inc., under license to Theratechnologies Inc.
THERA patient support® is a trademark of Theratechnologies Inc.
© 2023 Theratechnologies Inc. All rights reserved.
US-TRO-2200027-V2



Trogarzo®
(ibalizumab-uiyk)
Injection
200 mg/1.33 mL (150 mg/mL)