 **Trogarzo**[®]
(ibalizumab-uiyk)
Injection
200 mg/1.33 mL (150 mg/mL)

93% of TROGARZO[®] patients surveyed recommend TROGARZO[®] to other HIV-1 patients*
Discover more on page 4



Have you changed regimens more than 2 times in the past? Are you looking for a different option because your regimen is no longer working?

ASK YOUR HEALTHCARE PROVIDER ABOUT TROGARZO[®], THE FIRST LONG-ACTING HIV-1 TREATMENT

*Thirty patients completed a brief survey (online) regarding their experience with TROGARZO[®]. All surveys were completed from December, 2020 to March, 2021.

Please see Important Safety Information throughout and the enclosed full Prescribing Information for TROGARZO[®]

WHAT IS TROGARZO®?

TROGARZO® is a fully active medicine that's used in combination with other antiretroviral medicines to treat HIV-1 infection in adults who:



UNDETECTABLE:* A POSSIBILITY AGAIN

If treatment failure is part of your journey, you deserve to know about all of your treatment options, including TROGARZO® – HIV-1 treatment designed specifically for those with treatment failures.

Together, you and your healthcare provider can decide if TROGARZO® should be part of your treatment regimen to help keep your viral load in check.

ALL YOU HAVE TO DO IS ASK.

IMPORTANT SAFETY INFORMATION

Do not receive TROGARZO® if you have had an allergic reaction to TROGARZO® or any of the ingredients in TROGARZO®.

TROGARZO® can cause serious side effects, including:

- Allergic reactions. TROGARZO® can cause allergic reactions, including serious reactions, during and after infusion. Tell your healthcare provider or nurse, or get medical help right away if you get any of the following symptoms of an allergic reaction: trouble breathing, swelling in your throat, wheezing, chest pain, chest tightness, cough, hot flush, nausea or vomiting.
- Changes in your immune system (Immune Reconstitution Inflammatory Syndrome) can happen when you start taking HIV-1 medicines. Your immune system might get stronger and begin to fight infections that have been hidden in your body for a long time. Tell your healthcare provider right away if you start having new symptoms after receiving TROGARZO®.

Please see the back cover for complete safety details.

* Undetectable viral load is defined as fewer than 50 copies of HIV RNA per mL of blood.

LIGHTEN THE VIRAL LOAD

Did you know that when you become resistant to several HIV-1 medications, treatment guidelines note using a drug that works differently than others you may have tried?

TROGARZO® works differently. Unlike antiretroviral medicines that aim to prevent HIV-1 from multiplying inside the immune system's T-cells, TROGARZO® is an antibody bioengineered specifically to protect your T-cells from the virus. TROGARZO® binds to the T-cell's surface to block HIV-1 from entering.

THIS CAN HELP PROTECT THE T-CELL AND REDUCE OR "LIGHTEN" YOUR VIRAL LOAD.

In a clinical study, TROGARZO® was proven to have a significant and long-lasting impact on the amount of HIV-1 in the body, reducing or "lightening" the viral load.

83% OF PATIENTS WERE ABLE TO REDUCE THEIR HIV-1 VIRAL LOAD AFTER RECEIVING A SINGLE DOSE OF TROGARZO®.

Also, studies showed that patients who reached undetectable were able to remain undetectable over time.



TAKEN EVERY TWO WEEKS

Did you know that there is another option to taking daily pills?

TROGARZO® is not a daily pill. It's an intravenous (IV) infusion given over 15 to 30 minutes every 2 weeks.



Given once every 2 weeks



Appointments arranged by your personal Patient Care Coordinator



Given by a healthcare provider at home, an infusion center or healthcare provider's office

93% OF TROGARZO® PATIENTS SURVEYED RECOMMEND TROGARZO® TO OTHER HIV-1 PATIENTS*

93%

rated the TROGARZO® infusion process as “very convenient” or “neutral”.†

87%

were “very satisfied” with their TROGARZO® infusion experience.

87%

reported that having a 2-week touchpoint with their HCP was a positive experience.

- 80% reported that having a 2-week touchpoint with their HCP helped them adhere to their other HIV medications.

WHAT ARE THE BENEFITS OF AN IV INFUSION EVERY 2 WEEKS?

Infusion appointments allow you to keep in touch with your healthcare provider on a regular basis and they can be a great time to ask any questions you may have about your treatment or your health.

IMPORTANT SAFETY INFORMATION

Before you receive TROGARZO® (ibalizumab-uiyk), tell your healthcare provider about all of your medical conditions, including if you are:

- **Pregnant or plan to become pregnant.** It is not known if TROGARZO® may harm your unborn baby. Tell your healthcare provider if you become pregnant during treatment with TROGARZO®.
- **Breastfeeding or plan to breastfeed.** You should not breastfeed if you have HIV-1 because of the risk of passing HIV-1 to your baby. Do not breastfeed if you are receiving TROGARZO® as it is not known if TROGARZO® passes into breast milk. Talk with your healthcare provider about the best way to feed your baby during treatment with TROGARZO®.

Also tell your healthcare provider about all the medicines you take, including all prescription and over-the-counter medicines, vitamins, and herbal supplements.




* Thirty patients completed a brief survey (online) regarding their experience with TROGARZO®. All surveys were completed from December, 2020 to March, 2021.

† 93% of respondents rated the infusion process as “very convenient” (70%) or “neutral” (23%).

TROGARZO®, A POWERFUL, LONG-ACTING HIV-1 TREATMENT

Given once every 2 weeks, TROGARZO® is the first available long-acting antiretroviral against HIV-1 – a breakthrough in a class of its own.

How does the dosing of long-acting TROGARZO® make it different from pills?



If the level of medication in your body falls into the “red zone”, it means that the level may not be high enough to suppress your viral load and you can become resistant to your current medication.

TROGARZO® IS LONG-ACTING, WHICH MEANS THAT IT MAINTAINS A SUFFICIENT LEVEL OF MEDICATION IN THE BODY OVER ITS DOSING PERIOD OF 14 DAYS

A PROVEN TOLERABILITY & SAFETY PROFILE

TROGARZO® is generally well tolerated, with a low incidence of adverse reactions.

The most common side effects of TROGARZO® were diarrhea, dizziness, nausea, and rash. Most adverse reactions reported were mild to moderate in severity.

Did you know that some HIV-1 medications can cause interactions with other drugs which can reduce their efficacy or increase side effects?

TROGARZO® does not interact with other drugs, so you can take it with other medications you may be taking. Be sure to discuss other medications you may be taking with your healthcare provider before taking TROGARZO®.

IMPORTANT SAFETY INFORMATION

The most common side effects of TROGARZO® include diarrhea, dizziness, nausea, and rash. These are not all the possible side effects of TROGARZO®.

Do not receive TROGARZO® if you have had an allergic reaction to TROGARZO® or any of the ingredients in TROGARZO®.

Please see the back cover for complete safety details.

WHERE CAN I GET MY TROGARZO® INFUSION?

LOCATIONS AVAILABLE FOR INFUSIONS



INFUSION CENTER



YOUR HOME



YOUR HEALTHCARE PROVIDER'S OFFICE

Depending on your insurance coverage, your Patient Care Coordinator will arrange for your infusions to be given at a location of your choosing – at an infusion center, in your own home, or at your healthcare provider's office. If you decide over time that another location would be more convenient for you, simply let them know.

You can also reach them by phone to ask any questions you may have about your treatment.

87% of TROGARZO® patients surveyed were “very satisfied” with their TROGARZO® infusion experience

Discover more on page 4

WHAT TO EXPECT WITH YOUR INFUSIONS

Before your infusion

- Ask your Patient Care Coordinator if you should bring any documents (like insurance info) to your appointment.

During your infusion

- Someone will monitor your therapy during the 15-to-30-minute infusion.
- Take the opportunity to discuss HIV-related topics with your care provider.

After your infusion

- You'll also be monitored afterwards for a short period of time to ensure that your body tolerates your treatment well.

Please see the back cover for complete safety details.



“I've been HIV-positive for over 34 years. In the past 6½ years, my life has changed dramatically because I started TROGARZO® as part of a phase 2b study. Within 2 months, for the first time, I had undetectable viral loads. I didn't believe it at first, but as months went by, I started feeling more and more secure. And now we're talking 6½ years. I was able to restart my life again.”

This is the experience of one patient and may not be representative of all TROGARZO® patients.



SCAN HERE TO HEAR ABOUT THE EXPERIENCE OF NELSON VERGEL, A TROGARZO® PATIENT



MEET THERA patient support®

We're here to help make your journey as smooth as possible and answer any questions you may have.

Once TROGARZO® becomes a part of your regimen, you'll receive a welcome call from your dedicated Patient Care Coordinator.



Thinking about how to afford your treatment?

Don't worry, TROGARZO® is widely covered by all commercial and government insurance plans, including ADAP. Commercially insured patients may be eligible for co-pay assistance and may pay as little as \$0.

From appointment reminders to insurance coverage assistance, and anything in between, THERA patient support® is here to help.

LEARN MORE AT [THERAPATIENTSUPPORT.COM](https://therapatientssupport.com)



IS TROGARZO® RIGHT FOR YOU?

START THE CONVERSATION

Take this quiz to determine if TROGARZO® might be right for you:

Are you having any challenges taking your current HIV medications? Yes No

Do you skip or miss your HIV medication doses sometimes? Yes No

Have you changed your HIV medications in the past? Yes No

Are you taking your HIV medication and your viral load is still detectable? Yes No

Would you be open to including something other than a daily pill in your treatment regimen if given the choice? Yes No

Do you consider safety an important part of your next HIV medication choice? Yes No

If you answered "YES" to 4 out of 6 questions, you might be a potential TROGARZO® candidate.

AT YOUR NEXT VISIT WITH YOUR HEALTHCARE PROVIDER, ASK TO LEARN MORE ABOUT TROGARZO®!

When your HIV-1 treatment fails, it's important to stay on top of your treatment and know all of your options. Ask your healthcare provider if TROGARZO® should be part of your regimen to keep your viral load in check.

Because you deserve to know.

FOR MORE INFORMATION, VISIT [TROGARZO.COM](https://trogarzo.com)

You can view and download the discussion guide from the website! Remember to bring a copy with you to your next doctor's appointment.

Please see the next page for complete safety details.

SO, IS TROGARZO® RIGHT FOR YOU?

Have you changed regimens more than 2 times in the past?
Are you looking for a different option because your regimen
is no longer working?

LEARN MORE AT TROGARZO.COM

• THERA patient support®

If you have any questions, call the THERA patient support® program at

1-833-23 THERA (1-833-238-4372),

Monday-Friday, 8:30AM-8PM ET.

What is TROGARZO®?

TROGARZO® (ibalizumab-uiyk) is a prescription medicine that is used with other antiretroviral medicines to treat Human Immunodeficiency Virus-1 (HIV-1) infection in adults who:

- have received anti-HIV-1 regimens in the past, and
- have HIV-1 virus that is resistant to antiretroviral medicines, and
- who are failing their current antiretroviral therapy

It is not known if TROGARZO® is safe and effective in children.

IMPORTANT SAFETY INFORMATION

TROGARZO® can cause serious side effects, including:

- Allergic reactions. TROGARZO® can cause allergic reactions, including serious reactions, during and after infusion. Tell your healthcare provider or nurse, or get medical help right away if you

get any of the following symptoms of an allergic reaction: trouble breathing, swelling in your throat, wheezing, chest pain, chest tightness, cough, hot flush, nausea or vomiting.


- Changes in your immune system (Immune Reconstitution Inflammatory Syndrome) can happen when you start taking HIV-1 medicines. Your immune system might get stronger and begin to fight infections that have been hidden in your body for a long time. Tell your healthcare provider right away if you start having new symptoms after receiving TROGARZO®.

For more information or medical advice about side effects, ask your healthcare provider. You may report side effects to the FDA at 1-800-FDA-1088 or the THERA patient support® program at 1-833-238-4372.

Please see Important Safety Information throughout and the enclosed full Prescribing Information for TROGARZO®



TROGARZO® is a registered trademark of TaiMed Biologics Inc., under license to Theratechnologies Inc.
THERA patient support is a trademark of Theratechnologies Inc.
© 2021 Theratechnologies Inc. All rights reserved. 1024-01-09/21

 **Trogarzo®**
(ibalizumab-uiyk)
Injection
200 mg/1.33 mL (150 mg/mL)